

## Science News

from research organizations

### Carbon dioxide levels reflect COVID-19 risk

Research confirms value of measuring carbon dioxide to estimate infection risk

**Date:** April 7, 2021

**Source:** University of Colorado at Boulder

**Summary:** Tracking carbon dioxide levels indoors is an inexpensive and powerful way to monitor the risk of people getting COVID-19, according to new research. In any given indoor environment, when excess carbon dioxide levels double, the risk of transmission also roughly doubles, scientists report.

**Share:** [!\[\]\(d3102649f02e825ddb76dc3de0190154\_img.jpg\)](#) [!\[\]\(55ca3a38dbb940110628e54e3ea7505d\_img.jpg\)](#) [!\[\]\(1ad7b9dfa1e10e48660df5dd18a1b20c\_img.jpg\)](#) [!\[\]\(7b7f78f3b14c2b344e3d1b2a79a760c9\_img.jpg\)](#) [!\[\]\(ea85bb5cc605bcd0e585ecc5f09976f5\_img.jpg\)](#)

#### FULL STORY

Tracking carbon dioxide levels indoors is an inexpensive and powerful way to monitor the risk of people getting COVID-19, according to new research from the Cooperative Institute for Research in Environmental Sciences (CIRES) and the University of Colorado Boulder. In any given indoor environment, when excess CO<sub>2</sub> levels double, the risk of transmission also roughly doubles, two scientists reported this week in *Environmental Science & Technology Letters*.

The chemists relied on a simple fact already put to use by other researchers more than a decade ago: Infectious people exhale airborne viruses at the same time as they exhale carbon dioxide. That means CO<sub>2</sub> can serve as a "proxy" for the number of viruses in the air.

"You're never safe indoors sharing air with others, but you can reduce the risk," said Jose-Luis Jimenez, co-author of the new assessment, a CIRES Fellow and professor of chemistry at the University of Colorado Boulder.

"And CO<sub>2</sub> monitoring is really the only low-cost and practical option we have for monitoring," said Zhe Peng, a CIRES and chemistry researcher, and lead author of the new paper. "There is nothing else."

For many months, researchers around the world have been searching for a way to continually monitor COVID-19 infection risk indoors, whether in churches or bars, buses or hospitals. Some are developing instruments that can detect viruses in the air continually, to warn of a spike or to indicate relative safety. Others

tested existing laboratory-grade equipment that costs tens of thousands of dollars.

Jimenez and colleagues turned to commercially available carbon dioxide monitors, which can cost just a few hundred dollars. First, they confirmed in the laboratory that the detectors were accurate. Then, they created a mathematical "box model" of how an infected person exhales viruses and CO<sub>2</sub>, how others in the room inhaled and exhaled, and how the viruses and gas accumulate in the air of a room or are removed by ventilation. The model takes into consideration infection numbers in the local community, but it does not detail air flow through rooms -- that kind of modeling requires expensive, custom analysis for each room.

It's important to understand that there is no single CO<sub>2</sub> level at which a person can assume a shared indoor space is "safe," Peng emphasized. That's partly because activity matters: Are people in the room singing and talking loudly or exercising, or are they sitting quietly and reading or resting? A CO<sub>2</sub> level of 1,000 ppm, which is well above outside levels of about 400 ppm, could be relatively safe in a quiet library with masks but not in an active gym without masks.

But in each indoor space, the model can illuminate "relative" risk: If CO<sub>2</sub> levels in a gym drop from 2,800 to 1,000 ppm (~2,400 above background levels to 600), the risk of COVID-19 transmission drops, too, to one-quarter of the original risk. In the library, if an influx of people makes CO<sub>2</sub> jump from 800 to 1,600 (400 to 1,200 above background), COVID transmission risk triples.

In the new paper, Peng and Jimenez also shared a set of mathematical formulae and tools that experts in building systems and public health can use to pin down actual, not just relative, risk. But the most important conclusion is that to minimize risk, keep the CO<sub>2</sub> levels in all the spaces where we share air as low as practically possible.

"Wherever you are sharing air, the lower the CO<sub>2</sub>, the lower risk of infection," Jimenez said.

---

### Story Source:

Materials provided by **University of Colorado at Boulder**. *Note: Content may be edited for style and length.*

---

### Journal Reference:

1. Zhe Peng, Jose L. Jimenez. **Exhaled CO<sub>2</sub> as a COVID-19 Infection Risk Proxy for Different Indoor Environments and Activities**. *Environmental Science & Technology Letters*, 2021; DOI: 10.1021/acs.estlett.1c00183

---

### Cite This Page:

MLA	APA	Chicago
-----	-----	---------

---

University of Colorado at Boulder. "Carbon dioxide levels reflect COVID-19 risk: Research confirms value of measuring carbon dioxide to estimate infection risk." ScienceDaily. ScienceDaily, 7 April 2021. <[www.sciencedaily.com/releases/2021/04/210407143809.htm](https://www.sciencedaily.com/releases/2021/04/210407143809.htm)>.

---

## Featured Content

*from New Scientist*

Strongest evidence yet that MS is caused by Epstein-Barr virus

Jan. 13, 2022 — A huge study of US military personnel suggests almost all cases of multiple sclerosis are triggered by the common Epstein-Barr virus, meaning a vaccine could largely eradicate the condition.

Largest ever fish colony hosts 100 billion eggs under Antarctic ice

Jan. 13, 2022 — In the Weddell Sea near Antarctica, scientists have found the largest colony of fish nests in the world, covering 240 square kilometres.

Electric knee implants could help treat pain of osteoarthritis

Jan. 12, 2022 — A device that delivers electric current to the knees could help combat osteoarthritis, a painful condition caused by worn cartilage, after successful tests in rabbits.

Visit *New Scientist* for more global science stories >>>

## RELATED STORIES

---

Exploring the Relationship Between Nitrogen and Carbon Dioxide in Greenhouse Gas Emissions

Dec. 14, 2020 — An interdisciplinary study on a decade-long experiment (1997-2009) found that lower nitrogen levels in soil promoted release of carbon dioxide from soils under high levels of atmospheric carbon ...

Face Masks Unlikely to Cause Over-Exposure to Carbon Dioxide, Even in Patients With Lung Disease, Study Finds

Oct. 2, 2020 — New research findings contradict statements linking wearing face masks to carbon dioxide poisoning by trapping carbon dioxide. During the COVID-19 pandemic the wearing of face masks has become a ...

Artificial Photosynthesis Transforms Carbon Dioxide Into Liquefiable Fuels

May 22, 2019 — Chemists have successfully produced fuels using water, carbon dioxide and visible light through artificial photosynthesis. By converting carbon dioxide into more complex molecules like propane, green ...

Hunting Squid Slowed by Rising Carbon Levels

Mar. 21, 2018 — Scientists have found that high carbon dioxide levels cause squid to bungle attacks on their prey. Investigators said that the oceans absorb more than one-quarter of all the excess carbon dioxide ...

## FROM AROUND THE WEB

---

*ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.*

**Which air cleaners work best to remove aerosols that contain viruses?**

Robyn Schofield et al., TechXplore.com, 2021

**Researchers developing air quality sensors to detect COVID-19**

Janette Neuwahl Tannen et al., TechXplore.com, 2021

**A new program models airflows much faster than before, without a supercomputer**

Aalto University, TechXplore.com, 2021

## **Efficacy and safety following bosutinib dose reduction in patients with Philadelphia chromosome-positive leukemias**

Kota V et al. Leuk Res. 2021, Brought to you by Pfizer Medical affairs

## **Understanding “Atmosome”, the Personal Atmospheric Exposome: Comprehensive Approach**

Hari Bhimaraju et al., JMIR Biomedical Engineering, 2021

## **Versatile Graphene Underlies New COVID-Zapping Air Filter**

Author Leslie Mertz, IEEE PULSE, 2021

## **Scientific Communication in the Age of Coronavirus**

TrendMD Blog

## **Estimation of Viral Aerosol Emissions From Simulated Individuals With Asymptomatic to Moderate Coronavirus Disease 2019**

Michael Riediker et al., JAMA Network Open, 2020

---

Powered by **TREND MD**

## **Free Subscriptions**

---

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 Email Newsletters

 RSS Feeds

## **Follow Us**

---

Keep up to date with the latest news from ScienceDaily via social networks:

 Facebook

 Twitter

 LinkedIn

## **Have Feedback?**

---

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 Leave Feedback

 Contact Us

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners.

Content on this website is for information only. It is not intended to provide medical or other professional advice.

Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— CCPA: Do Not Sell My Information —